

A Narrative and Meta-Analytic Study of *in vivo* Efficiency of the Bioactive Compounds of Propolis in Tooth Decay

Yun Jin Kim¹, Yean Kong Yong², Muhammad Shahzad Aslam¹

ABSTRACT

Background: Propolis is one of the major components produced by honeybee. It is well known in different parts of the world such as Iran, Canada, Yemen, Czech Republic, Ethiopia, Bulgaria, Portugal, India, Turkey, Malaysia, the United States of America, Chile, Brazil, and Indonesia. The bioactive constituent of every type of propolis varies depending on the geographical location. Terpenoids, flavonoids, and polyphenol compounds were found to be common in all kinds of propolis. It possess numerous applications such as control of dental infections, plaque cleaning, treating gingivitis, exhibiting antimicrobial effect and treating radiation-induced oral mucositis and cariogenic infections in caries-active patients. **Methodology:** This study thus aimed to undertake a meta-analysis of the efficacy of bioactive compounds of propolis in tooth decay. A total of three *in vivo* studies were systematically reviewed, and two studies with a total of 300 pathogen-free female Wistar rats were included in the final meta-analysis. **Results:** The results were compared among three subcategories of smooth surface caries and sulcal caries (slight, moderate, and severe), supporting a statistically significant ($P = 0.006$) beneficial effect of using fractional propolis. **Conclusion:** Most of the included studies were preliminary, without blind study and lack of information about standard animal housing protocol. More *in vivo* and clinical trials of bioactive compounds of propolis should be encouraged in future.

Key words: Dental hygiene, *in vivo* studies, meta-analysis, propolis, tooth decay, traditional medicine

Yun Jin Kim¹, Yean Kong Yong², Muhammad Shahzad Aslam^{1*}

¹School of Traditional Chinese Medicine, Xiamen University Malaysia, MALAYSIA.

²Laboratory Centre, Xiamen University Malaysia, Jalan Sunsuria, Bandar Sunsuria, 43900 Sepang, Selangor Darul Ehsan, MALAYSIA.

Correspondence

Dr. Muhammad Shahzad Aslam,

School of Traditional Chinese Medicine, Xiamen University Malaysia, Jalan Sunsuria, Bandar Sunsuria, 43900 Sepang, Selangor Darul Ehsan, MALAYSIA.

Phone no : +60193009674

E-mail: aslam.shahzad@xmu.edu.my

DOI : 10.5530/phrev.2019.2.1

Article Available online

<http://www.phcogrev.com/v13/i26>

Copyright

© 2019 Phcog.Net. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.



INTRODUCTION

Dental caries, commonly known as tooth decay, is an infectious disease caused by dietary sugars, which results in loss of minerals and salts of the tooth and development of cavities.^[1] It has been found that sugar has a direct impact on tooth decay, but with the development of fluoride toothpaste, the prevalence of dental caries has significantly decreased despite increase in sugar consumption.^[2-4] A recent study also indicated that consumers with a high level of sugar consumption had higher dental caries prevalence.^[5] It is interesting to know that females are usually found to display greater prevalence rates as compared to males due to hormonal variations.^[6]

Apis are honeybees that give rise to well-known products such as beeswax, nectar, pollen, propolis, bee bread, bee brood, and royal jelly. Propolis is obtained from harvest resins from different botanical species. Honeybees collect them and bring them back to the colony to protect their hives from crack and cover them to defend against pathogens. It is famous due to its traditional medicinal properties such as antioxidant,^[7] anticancer especially oral cancer,^[8,9] antiviral,^[10] antidiabetic,^[11] anti-inflammatory,^[12] gingivitis,^[13] antiplaque,^[14] antimicrobial,^[15] immunomodulatory,^[16] giardiasis^[17] and estrogenic.^[18]

Propolis is very effective in dental care. A number of publications confirm its effectiveness in treatment

against plaque and gingivitis. Therefore, propolis can be used as mouthwash. Propolis mouthwash is found to be effective in Phase-2 clinical trial.^[19] Typified propolis (2%) was found more effective against mutans streptococci and lactobacilli than chlorhexidine.

Propolis mouthwash is found patient satisfaction (74%) than chlorhexidine (68%).^[20] The anticariogenic action of the commercially available propolis chewing gum with xylitol was observed in 30 healthy children aged 8–11 years with decayed, missing, and filled teeth and was found more effective than xylitol.^[21] The application of propolis in different dosage forms is mentioned in Table 1. A list of clinical trials related to oral health and their outcomes is mentioned in Table 2.

The color of propolis varies from region to region. For example, it is dark orange in the northern region, orange in the central coastal, dark brown in the central interior, and brown and dark brown in the southern region of Portugal.^[30] The color, aroma and flavor may vary in summer, autumn, and winter seasons.^[31] Poplar from Europe and Baccharis from Brazil are well-known propolis in the market, but there are some other kinds of propolis which depend on the geographical region. For example, Okinawan propolis is indigenous to Okinawa, Japan.^[32] Varia-

Cite this article: Jin KY, Kong YY, Aslam MS. A Narrative and Meta-Analytic Study of *in vivo* Efficiency of the Bioactive Compounds of Propolis in Tooth Decay. Pharmacogn Rev. 2019;13(26):28-35.

Table 1: Application of propolis in different dosage form

Dosage form	Methodology/formulation	Application/uses	Problem	Result	References
Mouthwash containing propolis	The alcohol-free mouthwash containing 5% w/v of Brazilian green propolis	Control of plaque and gingivitis	Current commercial brands contain alcohol	Alcohol-free mouthwash contains propolis could fight against plaque and gingivitis	[19]
Mouthwash containing propolis	2% typified propolis, mint flavor, polioxyethelers, sorbitol, blue color and water	Cariogenic infections in the caries-active patient	Dental infection	It is more effective than chlorhexidine	[20]
Propolis sachet	2 mL of AEP (13% solution)	Adjuvant therapy for asthma patient	Asthma	AEP is better than placebo	[22]
Propolis toothpaste	Silicate toothpaste with the extract from propolis	Plaque cleaning	Plaque	The toothpaste shows very good plaque-cleaning, plaque-inhibiting and anti-inflammatory effect	[23]
Chewing gum	The two chewing gums used in this study contained propolis 6.4% and xylitol 15% as the primary ingredient respectively	Natural anticarcinogenic agents	Cariogenic effect of <i>S. mutans</i>	It helps to reduce salivary <i>S. mutans</i> count when compared to xylitol gum	[21]
Mucoadhesive propolis gel	Brazilian green propolis, purified water, polysorbate 20, propylene glycol and hydroxypropyl methylcellulose	Radiation-induced oral mucositis	Mucositis is a common problem in cancer patient	Mucoadhesive propolis gel could be considered as a potential topical medication for preventing radiation-induced oral mucositis	[24]
Propolis - based chitosan varnish	EPE was ground into fine powder and 2.5 g of propolis was mixed with 25 mL of 80% aqueous ethanol in a test tube and shaken at 70°C for 30 min. After extraction, the mixture was centrifuged at 8000×g to obtain the supernatants. PCV were prepared by the addition of acetic acid to EPE 25%	Antimicrobial activity	Enhancement of propolis	The product is suitable for dental caries	[25]

S. mutans=*Streptococcus mutans*, AEP=Aqueous extract propolis, EPE=Ethanollic propolis extract, PCV=Propolis-based chitosan varnishes

Table 2: List of clinical trials related to oral health and their outcomes

Potential application	Type of clinical trial	Location	Number of patient	Outcome	References
Cavity disinfectants	Randomized clinical trial	Bapuji Dental College and Hospital, Davanagere	10	Propolis extracts can be used as a natural disinfecting agent	[26]
Control of plaque and gingivitis	Clinical trial phase-2	Faculty of Dentistry of Federa University of Minas Gerais, Brazil	25	It could be an alternative alcohol-free subject to further double-blind, randomized clinical trial	[19]
Cariogenic infections in a caries-active patient	Randomized, double-blind, placebo-controlled clinical trial	Dental Clinics at Bandeirante Anhanguera University - UNIBAN, Sao Paulo, Brazil	100	Effective against cariogenic infections when used 2% typified propolis, mint flavor, polioxyethelers, sorbitol, blue color and water	[20]
Dentinal hypersensitivity	A randomized, double-blind study	Department of Conservative Dentistry and Endodontics	156	Propolis was more effective than 5% potassium nitrate	[27]
CP and DMt2 receiving SRP	A masked, randomized clinical trial comparing with placebo	Internal Medicine Hospital, Mansoura University	50	400 mg daily propolis is a potentially viable adjunct to SRP that significantly reduces HbA1c, fasting plasma glucose, serum-(carboxymethyl) lysine levels and improves periodontal therapy outcomes in people with DMt2 and chronic periodontitis	[28]
Gingivitis	Randomized, double-blind, controlled clinical trial	N/A	42	2% typified propolis rinse was equivalent to a positive control rinse during a 21 days	[29]

CP=Chronic periodontitis, DMt2=Type 2 diabetes mellitus, SRP=Scaling and root planing, HbA1c=Hemoglobin A1c, N/A=Not available

Table 3: List of major type of propolis in a different part of the world

Origin	Pharmacological activity	<i>In vitro/in vivo</i> model	Bioactive compounds	Method identification of bioactive compounds	Result	References
Iranian propolis	Chemoprotection against gastric cancer	<i>In vivo</i>	Suberosin, tschimgin (bornyl p-hydroxybenzoate), tschimganin (bornyl vanillate), ferutinol (ferutinol p-hydroxybenzoate) and teferin (ferutinol vanillate)	NMR	A	[38,39]
Canadian propolis fractions	Anti-oxidant	<i>In vitro</i>	Chrysin, pinocembrin, palmitic acid, naringenin/pinobanksin, isopentyl caffeate, acacetin/caffeic acid	ESI-MS fingerprint	B	[40]
Yemen propolis	N/A	N/A	Triterpenoids, a-, p-amyryl, dammaradienyl acetates), n-alkenes, n-alkanes, n-alkanoic acids, long chain wax esters, n-alkanols and methyl n-alkanoates	GC-MS	C	[41]
Czech republic propolis	Antiviral	<i>In vitro</i>	Flavonoids, quercetin dihydrate, chrysin, pinocembrin Galangin, phenyl carboxylic acids, benzoic acid Cinnamic acid, caffeic acid, p-coumaric acid	HPLC	D	[10]
Ethiopian propolis	N/A	N/A	Triterpenoids such as a- and p-amyryns, a- and P-amyryl acetates, lupeol, a- and p-lupeyl acetates, n-alkanes, n-alkenes, long chain wax esters, methyl n-alkanoates	GC-MS	E	[42]
Bulgarian propolis	N/A	N/A	Pinocembrin, galangin, chrysin, tectochrysin, quercetin, isorhamnetin, kaempferol, 3,7-dihydroxy-5-methoxy flavanones and 2,5-dihydroxy-7-methoxy flavanones	HPLC, MS	F	[43]
Portugal propolis	Anti-inflammatory and antimicrobial	<i>In vitro</i>	N/A	N/A	G	[44]
Bulgarian propolis	N/A	N/A	Dihydrocaffeic acid, dihydroferulic acid, dihydroxyacetophenone, hydroxymethoxyacetophenone, p-phenethyl alcohol, benzyl alcohol pinobanksin, pinostrobin, dimethyl kaempferol	Capillary GC-MS	H	[45]
Southern portugal	Antioxidant activity	<i>In vitro</i>	Polyphenol and flavonoids	UV	I	[46]
Indian propolis	Antioxidant activity	<i>In vitro</i>	Pinocembrin and galangin	NMR	J	[47]
Portuguese propolis	Antioxidant and chemopreventive	<i>In vitro</i>	Ellagic acid, luteolin, a dimethoxylated Flavonol, dihydroxy-dimethoxyflavone, chrysoeriol-methyl ether, quercetin and kaempferol derivatives, dihydroflavones such as dihydroflavonols such as pinobanksin-3-O-pentenoate and pinobanksin-3-O-hexanoate	LC-DAD-ESI-MSn	K	[48,49]
Turkish propolis	Antioxidant and anti-cancer	<i>In vitro</i>	Polyphenols	UV	L	[50]
MP	Anti-oxidant and cardioprotective	<i>In vivo</i>	Flavonoids and polyphenols	UV	M	[51]
Indonesian propolis	Antioxidant	<i>In vitro</i>	Propolin d, Propolin c, Propolin f Propolin g, 5-pentadecylresorcinol, 5-heptadecylresorcinol, 5-(8z, 11z heptadecadienyl)-resorcinol, 5-(11z heptadecenyl)-resorcinol	NMR	N	[52]

A=Iranian propolis is beneficial in N-methyl-N-nitro-N-nitrosoguanidine induced gastric cancer at 100 ug/mL through inhibition of cell proliferation and apoptosis induction, B=Ethanol extract of Canadian propolis fractions showed higher polyphenol and flavonoid concentrations and higher antioxidant capacity as compared to commercial propolis and water extracts, C=Bioactive compounds indicates its pharmacological potential, D=Propolis extracts might be suitable for topical application against herpes infection, E=The difference among chemical composition of propolis in all parts of the world is due to diverse environmental source vegetation, F=The chemical composition of Bulgarian propolis is different from USSR due to geographical difference as bees collect propolis from resinous tree buds, G=Hydroalcoholic extract showed promising anti-microbial and anti-inflammatory activities, H=North and South Bulgaria have identified fifteen phenolic acids, I=There could be the possibility of difference in quality of propolis due to different zones in the same area, J=Galangin possess the highest anti-oxidant activity as compared to pinocembrin, K=Propolis exhibit anti-cancer activity on primary cultured cancerous renal cells, L=Ethanol extract of Turkish propolis possess good antioxidant and anticancer properties, M=MP exhibits cardioprotective activity against ISO-induced oxidative stress through its direct cytotoxic radical-scavenging activities, N=All prenylflavonones demonstrated significant radical scavenging activity against diphenylpicrylhydrazyl radicals, MP=Malaysian propolis, N/A=Not available

tion of bioactive compounds is huge from region to region and among different countries. Even within one country, there exists biochemical variation. For example, Brazil propolis has classified into 13 groups.^[33] A number of bioactive compounds have been isolated from Brazil propolis such as 3-prenyl-4-dihydrocinnamoxycinnamic acid, 2,2-dimethyl-6-carboxyethenyl-2-H1benzopyran, 3-prenyl-4-hydroxycinnamic acid, 2,2-dimethyl-6-carboxyethenyl-2-H1benzopyrane, 3,5-diprenyl-4-hy-

droxycinnamic acid, 2,2-dimethyl-6-carboxyethenyl-8-prenyl-2-H-1-benzopyran, labdanetype diterpenes, and prenylated chromane derivative.^[34-37] Table 3 enlists propolis of different regions with their chemical constituents and pharmacological activities.

There is an increase in the prevalence of tooth decay among preschool and schoolchildren in different countries such as India^[53] and Saudi Arabia^[54] and tropical countries such as Vietnam, Cambodia, Indonesia and

Myanmar.^[55] It is essential to explore the role of bioactive compounds in biomedicine. The selection of propolis is based on available literature as potential treatment options. In 2014, a meta-analysis has reported statistically insignificant evidence of an effect on reducing dental plaque using propolis. The results have shown an improvement in reducing dental plaque, but the overall results were not significant.^[56] Till now, no investigation has made an effort to understand the relationship of potential bioactive compounds of propolis and their preclinical or clinical effect on oral health. Most of the meta-analyses were based on the outcome of propolis in a different clinical trial. There is a huge variation among bioactive compounds of propolis even within same countries which no one accounts in the previous meta-analysis. The role of bioactive compounds and their *in vivo* efficacy are unclear. Hence, this meta-analysis aimed to investigate the efficacy of bioactive compounds of propolis for preventing tooth decay in animals.

METHODOLOGY

A systematic literature search was conducted in accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines using keywords propolis and/or bioactive compounds and/or clinical trial and/or *in vivo* studies and/or animal studies. A preliminary search on the PubMed, PubMed Central, CNKI, Scopus, Web of Science, Google Scholar, and PsycINFO databases yielded 15,537 papers published in English between January 1, 2000, and December 31, 2018. Inclusion and exclusion criteria were based on the study design, participants,

intervention, outcome criteria as mentioned in Table 4. Studies included in the systematic review focused on identified bioactive compounds mentioned in Table 5. Characteristics of the studies included are mentioned in Table 6.

Methodological quality (MQ) was studied using a wide-ranging 12-item assessment as mentioned in Figure 1. MQ scores were based on yes (1) or no (1). Usage of systematic reviews (SRs) of experimental animal studies is not yet a common practice, but awareness of the merits of conducting such SRs is steadily increasing. Systematic Review Centre for Laboratory Animal Experimentation's (SYRCLE) riskofbias (RoB) tool for animal studies is used in this study as mentioned in Table 7. The SYRCLE's RoB tool is an adapted version of the Cochrane RoB tool.^[57]

Statistical analysis

Meta-analysis was conducted using the Review Manager 5.3 software. The summary measures were reported as odds ratios or as a standard mean difference with 95% confidence intervals. The presence of heterogeneity among trials was assessed using the Chi-square test, and the extent of inconsistency was measured by I^2 statistics.^[58]

RESULTS

After screening around 15,573 articles, only two full-text papers were selected for further review. The abstraction process and reasons for exclusion are detailed in Figure 2. Table 5 summarizes the details of the studies reviewed. Two studies^[59,60] were random *in vivo* studies. One

Table 4: Study design; participants; intervention; outcome criteria

	Inclusion criteria	Exclusion criteria
Study design	Mixed method	All review articles, irrelevant articles, articles on chemical properties and optimization, <i>in vitro</i> activities, exclude tentative identification of bioactive compounds with <i>in vitro</i> activities, exclude <i>in vivo</i> activities with no information on bioactive compounds, exclude <i>in vivo</i> activities on different pharmaceutical preparation, exclude paper that uses propolis in combination with other agent, exclude studies due to absence in scoring rat dental caries by Keyes' method, exclude <i>in vivo</i> activities other than caries development/dental caries development
Participants	Pathogen-free female wistar rats	
Intervention	Propolis extract and their fractions delivered to pathogen-free female wistar rats/pup, infected with <i>S. sobrinus</i> and <i>S. mutans</i>	
Outcomes	Role of the bioactive composition of propolis extract/fractions in caries development	

S. sobrinus=*Streptococcus sobrinus*, *S. mutans*=*Streptococcus mutans*

Table 5: Studies included in the systematic review

Author, year	Country	Sample size in each group (n)	Study groups	Identified bioactive compounds	Identification method	Conclusions
Duarte, 2006	Brazil	12	Crude propolis (EEP), hexane fraction (EEH), the vehicle control ethanol 80%	Benzenepropanoic acid, methyl ester, trans-caryophyllene, a-humulene, nerolidol, myristic acid, 1,2-benzenedicarboxylic acid, bis (2-methylpropyl) ester, palmitic acid, methyl ester, hexadecanoic acid, methyl ester, linoleic acid, methyl ester, oleic acid, methyl ester, stearic acid, methyl ester, behenic acid, methyl ester, 1,2-benzenedicarboxylic acid, bis (2-ethylhexyl) ester	GC-MS	Oleic acid, linoleic acid, palmitic acid and stearic acid found in nonpolar bioactive hexane fraction of propolis and was able to reduce the incidence and severity of sulcal surface caries (P<0.05)
Silva, 2013	Brazil	13	NV (800 µg/mL), 250 ppm fluoride and vehicle control (20% ethanol, v/v)	NV	GC-MS	NV containing fraction isolated from Brazilian red propolis as effective as fluoride in reducing the development of carious lesions <i>in vivo</i>

NV=Neovestitol-vestitol

study on the *in vitro* and *in vivo* effects of isolated fractions of Brazilian propolis on caries development was removed due to the absence of scoring rat dental caries by Keyes' method.^[61] SYRCLE's RoB tool for animal studies is used in this study as mentioned in Table 7. In both of the studies, there was lack of information about the conditions of the animal house, animal loss during the study, attempt to blind the researcher, validity and reliability measures, and missing data reported as mentioned in Figure 1. The hexane fraction of Brazilian propolis (type 6) was compared with the control ethanol 80% in Duarte *et al.* experimentation, whereas neovestitol-vestitol containing fraction isolated from Brazilian red propolis was compared with the control (fluoride) in Bueno-Silva *et al.* experiment. The comparison was made between smooth surface caries and sulcal caries. The subgroups were divided into slight, moderate and severe

according to Keyes' method. In the assessment of potential publication bias, visual inspection of the funnel plot in smooth surface caries and sulcal caries revealed a roughly symmetrical distribution of studies as shown in Figures 3 and 4, respectively. The overall effect in smooth surface caries and sulcal caries was found statistically significant ($P = 0.006$). Heterogeneity test in general on smooth surface caries ($I^2 = 36\%$) and sulcal caries ($I^2 = 65\%$) differed comparatively. Therefore, smooth surface caries showed low heterogeneity as compared to sulcal caries, whereas in both cases, there was no heterogeneity within the groups (i.e., slight, moderate, and severe). Chi-square test, df values and I^2 statistics of every individual group are mentioned in Figures 5 and 6.

DISCUSSION

The results of this meta-analysis suggest that fractional propolis is beneficial in preventing tooth decay. The reason for low heterogeneity in smooth surface caries is because of less potential for plaque attachment as compared to sulcal caries. Sulcal caries is the potential area between the tooth and gingival tissue. Hence, it is the target site for tooth decay.

The studies examined in this article are in concordance with the fact that propolis improves the oral health; however, these finding lacks validity because of less available studies. Moreover, there is a need to isolate identified bioactive compounds and evaluate their preclinical and clinical trial.

From the literature, it was also found that the identified bioactive compounds in propolis may possess antimicrobial properties. Nerolidol present in green tea possesses antimicrobial activity.^[62,63] Trans-caryophyllene possesses antimicrobial activity against cariogenic bacteria.^[64] α -Humulene exhibits antibacterial activity against *Streptococcus mutans*.^[65] Myristic acid and palmitic acid exhibit antimicrobial activity against oral microorganisms such as *S. mutans*, *S. gordonii* and *S. sanguinis*.^[66] Behenic acid has been used with dental fluoride foam as an emulsion stabilizer.^[67] Neovestitol and vestitol exhibit anti-inflammatory and antimicrobial properties against *Actinomyces naeslundii*, *S. mutans*, *Staphylococcus aureus*, and *S. sobrinus*.^[68-70]




Figure 1: Methodological quality assessment of the two studies included in the meta-analysis (0 = no/not reported, 1 = yes)

Table 6: Study characteristics

Author, year	Study design	Intervention	Comparison group (s)	Statistical analysis	Software	Improved outcomes
Duarte, 2006	<i>In vivo</i> animal experimentation	Effect of chemical composition of propolis extract/fractions delivered to pathogen free female wistar rats infected with <i>S. sobrinus</i>	Crude propolis (EEP), hexane fraction (EEH), the vehicle control ethanol 80%	ANOVA, Tukey-Kramer HSD test	JMP version 3.1 software	Significant improvement when compared with the control group however unable to find substantial evidence of the role of identified bioactive compounds due to limitation as it require further investigation
Silva, 2013	<i>In vivo</i> animal experimentation	Effect of chemical composition of propolis extract/fractions delivered to pathogen free female wistar rats/pup infected with <i>S. mutans</i>	NV (800 ng/mL), 250 ppm fluoride and vehicle control (20% ethanol, v/v)	ANOVA, Tukey-Kramer HSD test for all pairs	BioEst version 5.0	Significant improvement when compared with the control group however unable to find substantial evidence of the role of identified bioactive compounds due to limitation as it require further investigation

Table 7: Results of SYRCLE's risk of bias tool for animal studies

Study (author, year)	Sequence generation	Baseline characteristics	Allocation concealment	Random housing	Blinding	Random outcome assessment	Incomplete outcome data	Selective outcome reporting	Other sources of bias
Duarte, 2006	Unclear	Yes	No	Unclear	Yes	Unclear	Unclear	Yes	Yes
Silva, 2013	Unclear	Yes	No	Unclear	Yes	Unclear	Unclear	Yes	Yes




Figure 2: Summary of search strategy




Figure 4: Forest plot in sulcal caries




Figure 5: Funnel plot showing overall standardized mean difference in smooth surface caries




Figure 3: Forest plot in smooth surface caries




Figure 6: Funnel plot showing overall standardized mean difference in sulcal caries

On the other hand, the limitations of the present meta-analysis should be debatable. First, the outcomes of this study should be inferred as carefully optimistic because of the unavailability of data. Most of the studies on propolis are either on crude or extract. Second, there is a lack of literature on isolated bioactive compounds on propolis and moreover, most of the studies are limited to *in vitro* or *in vivo* studies. Animal studies need to be more robust and well defined. In general, there are a number of phytochemical studies on propolis, but these studies cannot proceed further for the isolation and evaluation of identified compounds for a clinical trial in tooth decay. From the literature, it was also found that there is no clinical report on identified compounds other than dental caries.

CONCLUSION

Existing evidence proposes that fractional propolis is beneficial in preventing tooth decay. However, there are few studies currently available that restrict the currently available evidence. No studies are available that have made an effort to isolate bioactive compounds and test those bioactive compounds either on animal model or under clinical trial. Moreover, there is a need for research in the standardization of propolis which results in variability in their bioactive compounds. The variation in bioactive compounds results in inconsistent results which is the reason for insignificant statistical results from the previous meta-analysis of a clinical trial performed on propolis. There is a need to isolate identified compounds and evaluate their clinical potential to validate the results and determine the final conclusion.

ACKNOWLEDGMENT

This research was supported by Xiamen University Malaysia. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

ABBREVIATIONS

AEP: Aqueous extract propolis; **EPE:** Ethanolic propolis extract; **PCV:** Propolis based chitosan varnishes; **CP:** Chronic periodonti-tis; **Dmt2:** Type 2 diabetes mellitus; **SRP:** Scaling and root planning; **HbA1c:** Hemoglobin A1c; **NMR:** Nuclear magnetic resonance spectroscopy; **ESI-MS:** Electrospray ionization-Mass spectroscopy; **HPLC:** High Performance Liquid Chromatography; **GCMS:** Gas chromatography-mass spectrometry; **UV:** Ultraviolet Spectroscopy; **LC-DAD:** Liquid chromatography with diode array detector; **N/A:** Not available.



REFERENCES

- Lamont RJ, Eglund PG. Dental Caries. In: Molecular Medical Microbiology. Elsevier, 2015:945-55.
- Bernabé E, Vehkalahti MM, Sheiham A, Lundqvist A, Suominen AL. The Shape of the Dose-Response Relationship between Sugars and Caries in Adults. *J Dent Res* 2016;95:167-72.
- Woodward M, Walker AR. Sugar consumption and dental caries: evidence from 90 countries. *Br Dent J* 1994;176:297-302.
- Touger-Decker R, van Loveren C. Sugars And Dental Caries. *J Public Health Dent* 1944;4:7.
- Peres MA, Sheiham A, Liu P, Demarco FF, Silva AER, Assunção MC, et al. Sugar Consumption and Changes in Dental Caries from Childhood to Adolescence. *J Dent Res* 2016;95:388-94.
- Lukacs JR, Largaespada LL. Explaining sex differences in dental caries prevalence: Saliva, hormones, and 'life-history' etiologies. *Am J Hum Biol* 2006. doi:10.1002/ajhb.20530.
- Pazin WM, Mônico L da M, Egea Soares AE, Miguel FG, Berretta AA, Ito AS. Antioxidant activities of three stingless bee propolis and green propolis types. *J Apic Res* 2017;56:40-9.

- Sawicka D, Car H, Borawska MH, Niklinski J. The anticancer activity of propolis. *Folia Histochem Cytobiol* 2012;50:2537.
- Vagish Kumar LS. Propolis in dentistry and oral cancer management. *North American Journal of Medical Sciences*. 2014;6:11-20.
- Schnitzler P, Neuner A, Nolkemper S, Zundel C, Nowack H, Sensch KH, et al. Antiviral activity and mode of action of propolis extracts and selected compounds. *Phyther Res* 2010;24(SUPPL. 1). doi:10.1002/ptr.2868.
- Kang LJ, Lee HB, Bae HJ, Lee SG. Antidiabetic effect of propolis: Reduction of expression of glucose-6-phosphatase through inhibition of Y279 and Y216 autophosphorylation of GSK-3 α/β in HepG2 cells. *Phyther Res* 2010;24:155461.
- Park EH, Kim SH, Park SS. Anti-inflammatory activity of propolis. *Arch Pharm Res* 1996;19:33741.
- Kasuma N. Propolis Toothpaste as The Initial Therapy of Mild Gingivitis. *J Sains Farm Klin* 2014;1:89-94.
- Bapat S, Asawa K, Tak M, Chaturvedi P, Gupta V, George P, et al. The antiplaque efficacy of propolis-based herbal toothpaste: A crossover clinical study. *J Nat Sci Biol Med* 2015;6:364.
- Castro ML, Vilela WR, Zauli RC, Ikegaki M, Rehder VLG, Foglio MA, et al. Bioassay guided purification of the antimicrobial fraction of a Brazilian propolis from Bahia state. *BMC Complement Altern Med* 2009;9. doi:10.1186/1472-6882-9-25.
- Rindt IK, Spinu M, Niculae M. The Immunomodulatory Effect of Propolis : a Review. *Lucr Stiintifice Med Vet* 2009;2009;42:35.
- Abdel-Fattah NS, Nada OH. Effect of propolis versus metronidazole and their combined use in treatment of acute experimental giardiasis. *J Egypt Soc Parasitol* 2007;37(2 Suppl):691710.
- Okamoto Y, Tobe T, Ueda K, Takada T, Kojima N. Oral administration of Brazilian propolis exerts estrogenic effect in ovariectomized rats. *J Toxicol Sci* 2015;40:235-42.
- Santos VR, Pereira EMR, Da Silva JLDC, Silva FF, De Luca MP, Ferreira EFE, et al. Clinical evidence of the efficacy of a mouthwash containing propolis for the control of plaque and gingivitis: A phase II study. *Evidencebased Complement Altern Med* 2011. doi:10.1155/2011/750249.
- Netto CA, Marcucci MC, Paulino N, Anido-Anido A, Amore R, de Mendonca S, et al. Effects of typhoid propolis on mutans streptococci and lactobacilli: a randomized clinical trial. *Braz Dent Sci* 2013;16:31-6.
- Tulsani S, Chikkanarasaiah N, Siddaiah S, Krishnamurthy N. The effect of Propolis and Xylitol chewing gums on salivary Streptococcus mutans count: A clinical trial. *Indian J Dent Res* 2014;25:737.
- Khayyal MT, El-Ghazaly MA, El-Khatib AS, Hatem AM, De Vries PJ, El-Shafei S, et al. A clinical pharmacological study of the potential beneficial effects of a propolis food product as an adjuvant in asthmatic patients. *Fundam Clin Pharmacol* 2003;17:93102.
- Botushanov PI, Grigorov GI, Aleksandrov G a. A clinical study of a silicate toothpaste with extract from propolis. *Folia Med (Plovdiv)* 2001;43:28-30.
- Noronha VRAS, Araujo GS, Gomes RT, Iwanaga SH, Barbosa MC, Abdo EN, et al. Mucoadhesive propolis gel for prevention of radiation-induced oral mucositis. *Curr Clin Pharmacol* 2014;9:359-64.
- Franca JR, De Luca MP, Ribeiro TG, Castilho RO, Moreira AN, Santos VR, et al. Propolis -based chitosan varnish: Drug delivery, controlled release and antimicrobial activity against oral pathogen bacteria. *BMC Complement Altern Med* 2014;14:111.
- Prabhakar A, Karuna Y, Yavagal C, Deepak B. Cavity disinfection in minimally invasive dentistry -comparative evaluation of Aloe vera and propolis: A randomized clinical trial. *Contemp Clin Dent* 2015;6:24.
- Purra A, Mushtaq M, Acharya S, Saraswati V. A comparative evaluation of propolis and 5.0% potassium nitrate as a dentine desensitizer: A clinical study. *J Indian Soc Periodontol* 2014;18:466.
- El-Sharkawy HM, Anees MM, Van Dyke TE. Propolis Improves Periodontal Status and Glycemic Control in Patients With Type 2 Diabetes Mellitus and Chronic Periodontitis: A Randomized Clinical Trial. *J Periodontol* 2016;87:1418-26.
- Bretz WA, Paulino N, Nör JE, Moreira A. The Effectiveness of Propolis on Gingivitis: A Randomized Controlled Trial. *J Altern Complement Med* 2014;20:943-8.
- Falcão SI, Freire C, Vilas-Boas M. A proposal for physicochemical standards and antioxidant activity of Portuguese propolis. *JAOCS, J Am Oil Chem Soc* 2013;90:172941.
- Figueiredo FJB, Dias-Souza MV, Nascimento EA, De Lima LRP. Physicochemical characterization and flavonoid contents of artisanal Brazilian green propolis. *Int J Pharm Pharm Sci* 2015;7:64-8.
- Kumazawa S, Nakamura J, Murase M, Miyagawa M, Ahn MR, Fukumoto S. Plant origin of Okinawan propolis: honeybee behavior observation and phytochemical analysis. *Naturwissenschaften* 2008;95:781-6.
- Rufatto LC, dos Santos DA, Marinho F, Henriques JAP, Roesch Ely M, Moura S. Red propolis: Chemical composition and pharmacological activity. *Asian Pac J Trop Biomed* 2017;7:59-8.
- Banskota AH, Tezuka Y, Prasain JK, Matsushige K, Saiki I, Kadota S. Chemical constituents of Brazilian propolis and their cytotoxic activities. *J Nat Prod* 1998;61:896900.
- Banskota AH, Tezuka Y, Adnyana IK, Ishii E, Midorikawa K, Matsushige K, et al.

- Hepatoprotective and anti-Helicobacter pylori activities of constituents from Brazilian propolis. *Phytomedicine* 2001;8:16-23.
36. Aga H, Shibuya T, Sugimoto T, Kurimoto M, Nakajima S. Isolation and Identification of Antimicrobial Compounds in Brazilian Propolis. *Biosci Biotechnol Biochem* 1994. doi:10.1271/bbb.58.945.
 37. Marcucci MC, Ferreres F, García-Viguera C, Bankova VS, De Castro SL, Dantas AP, et al. Phenolic compounds from Brazilian propolis with pharmacological activities. *J Ethnopharmacol* 2001. doi:10.1016/S03788741(00)003263.
 38. Alizadeh AM, Afrouzan H, Dinparast-Djadid N, Sawaya ACHF, Azizian S, Hemmati HR, et al. Chemoprotection of MNNG-initiated gastric cancer in rats using Iranian propolis. *Arch Iran Med* 2015;18:18-23.
 39. Trusheva B, Todorov I, Ninova M, Najdenski H, Daneshmand A, Bankova V. Antibacterial mono- and sesquiterpene esters of benzoic acids from Iranian propolis. *Chem Cent J* 2010;4:1-4.
 40. Cottica SM, Sabik H, Antoine C, Fortin J, Graveline N, Visentainer J V., et al. Characterization of Canadian propolis fractions obtained from two-step sequential extraction. *LWT Food Sci Technol* 2015;60:60914.
 41. Al-Ghamdi AA, Bayaqoob NIM, Rushdi AI, Alattal Y, Simoneit BRT, El-Mubarak AH, et al. Chemical compositions and characteristics of organic compounds in propolis from Yemen. *Saudi J Biol Sci* 2017;24:1094103.
 42. Rushdi AI, Adgaba N, Bayaqoob NIM, Alkhazim A, Simoneit BIT, El-Mubarak AH, et al. Characteristics and chemical compositions of propolis from Ethiopia. *Springerplus* 2014;3:1-9.
 43. Bankova VS, Popov SS, Marekov NL. A study on flavonoids of propolis. *J Nat Prod* 1983;46:471-4.
 44. Silva JC, Rodrigues S, Feás X, Estevinho LM. Antimicrobial activity, phenolic profile and role in the inflammation of propolis. *Food Chem Toxicol* 2012;50:1790-5.
 45. Bankova V, Dylgierov AI, Popov S, M N. A GC / MS Study of the Propolis Phenolic Constituents. 2015;151:14751.
 46. Miguel MG, Nunes S, Dandlen SA, Cavaco AM, Antunes MD. Phenols and antioxidant activity of hydroalcoholic extracts of propolis from Algarve, South of Portugal. *Food Chem Toxicol* 2010;48:3418-23.
 47. Laskar RA, Sk I, Roy N, Begum NA. Antioxidant activity of Indian propolis and its chemical constituents. *Food Chem* 2010;122:2337.
 48. Valente MJ, Baltazar AF, Henrique R, Estevinho L, Carvalho M. Biological activities of Portuguese propolis: Protection against free radical-induced erythrocyte damage and inhibition of human renal cancer cell growth *in vitro*. *Food Chem Toxicol* 2011;49:86-92.
 49. Falcão SI, Vale N, Gomes P, Domingues MRM, Freire C, Cardoso SM, et al. Phenolic profiling of Portuguese propolis by LC-MS spectrometry: Uncommon propolis rich in flavonoid glycosides. *Phytochem Anal* 2013;24:30918.
 50. Turan I, Demir S, Misir S, Kilinc K, Mentese A, Aliyazicioglu Y, et al. Cytotoxic effect of Turkish propolis on liver, colon, breast, cervix and prostate cancer cell lines. *Trop J Pharm Res* 2015;14:777-82.
 51. Ahmed R, Tanvir EM, Hossen MS, Afroz R, Ahmmed I, Rumpa NEN, et al. Antioxidant properties and cardioprotective mechanism of Malaysian propolis in rats. *Evidence-based Complement Altern Med* 2017;2017. doi:10.1155/2017/5370545.
 52. Trusheva B, Popova M, Koendhori EB, Tsvetkova I, Naydenski C, Bankova V. Indonesian propolis: Chemical composition, biological activity and botanical origin. *Nat Prod Res* 2011;25:60613.
 53. Aranganal P, Mahadev SK, Jayaprakash J. Prevalence of Dental Caries among School Children in Chennai, Based on ICDAS II. *J Clin Diagn Res* 2016;10:ZC0912.
 54. Al-Meedani LA, Al-Dlaigan YH. Prevalence of dental caries and associated social risk factors among preschool children in Riyadh, Saudi Arabia. *Pakistan J Med Sci* 1969;32. doi:10.12669/pjms.322.9439.
 55. Bourgeois DM, Llodra JC. Global burden of dental condition among children in nine countries participating in an international oral health promotion programme, 2012-2013. *Int Dent J* 2014;64:27-34.
 56. Hwu YJ, Lin FY. Effectiveness of propolis on oral health: a meta-analysis. *J Nurs Res* 2014;22:221-9.
 57. Chen KY, Ma B, Wang YN, Chen CH, Zhao YQ, Zheng JX, et al. SYRCL's risk of bias tool for animal studies. *Chinese J EvidenceBased Med* 2014. doi:10.7507/16722531.20140206.
 58. Cochrane TC. Review Manager (RevMan) 5.3. Copenhagen Nord Cochrane Cent 2008.
 59. Bueno-Silva B, Koo H, Falsetta ML, Alencar SM, Ikegaki M, Rosalen PL. Effect of neovestitol-vestitol containing Brazilian red propolis on accumulation of biofilm *in vitro* and development of dental caries *in vivo*. *Biofouling* 2013;29:1233-42.
 60. Duarte S, Rosalen PL, Hayacibara MF, Cury JA, Bowen WH, Marquis RE, et al. The influence of a novel propolis on mutans streptococci biofilms and caries development in rats. *Arch Oral Biol* 2006. doi:10.1016/j.archoralbio.2005.06.002.
 61. Hayacibara MF, Koo H, Rosalen PL, Duarte S, Franco EM, Bowen WH, et al. *In vitro* and *in vivo* effects of isolated fractions of Brazilian propolis on caries development. *J Ethnopharmacol* 2005;101:110-5.
 62. Perumalla AVS, Hettiarachchy NS. Green tea and grape seed extracts — Potential applications in food safety and quality. *Food Res Int* 2011;44:827-39.
 63. Kubo I, Muroi H, Himejima M. Antimicrobial activity of green tea flavor components and their combination effects. *J Agric Food Chem* 1992;40:245-8.
 64. Aguiar GP, Carvalho CE, Dias HJ, Reis EB, Martins MHG, Wakabayashi KAL, et al. Antimicrobial activity of selected essential oils against cariogenic bacteria. *Nat Prod Res* 2013;27:1668-72.
 65. Kubo I, Muroi H, Kubo A. Antibacterial activity of long-chain alcohols against *Streptococcus mutans*. *J Agric Food Chem* 1993;41:2447-50.
 66. Huang CB, Alimova Y, Myers TM, Ebersole JL. Short- and medium-chain fatty acids exhibit antimicrobial activity for oral microorganisms. *Arch Oral Biol* 2011. doi:10.1016/j.archoralbio.2011.01.011.
 67. Edwin I. Stoltz. Dental fluoride foam. 2001. <https://patents.google.com/patent/US6251369B1/en> [Last accessed on 2018 Dec 27].
 68. Franchin M, Colón DF, da Cunha MG, Castanheira FVS, Saraiva ALL, Bueno-Silva B, et al. Neovestitol, an isoflavonoid isolated from Brazilian red propolis, reduces acute and chronic inflammation: involvement of nitric oxide and IL6. *Sci Rep* 2016;6:36401.
 69. Franchin M, Colón DF, Castanheira FVS, da Cunha MG, Bueno-Silva B, Alencar SM, et al. Vestitol Isolated from Brazilian Red Propolis Inhibits Neutrophils Migration in the Inflammatory Process: Elucidation of the Mechanism of Action. *J Nat Prod* 2016;79:954-60.
 70. Bueno-Silva B, Alencar SM, Koo H, Ikegaki M, Silva GVJ, Napimoga MH, et al. Anti-Inflammatory and Antimicrobial Evaluation of Neovestitol and Vestitol Isolated from Brazilian Red Propolis. *J Agric Food Chem* 2013;61:4546-50.

GRAPHICAL ABSTRACT



Assistant Prof. Dr. Yeon Kong Yong; Deputy Director, Laboratory Center, Xiamen University Malaysia.

SUMMARY

Propolis produced by honeybee is geographic centric and difficult to standardize. The propose of current article is to evaluate the qualitative synthesis and meta-analysis of *in-vivo* studies of propolis in tooth decay.

ABOUT AUTHORS



Associate Prof. Dr. Yun Jin Kim; School of Traditional Chinese Medicine, Xiamen University Malaysia.



Dr. Muhammad Shahzad Aslam; Lecturer, School of Traditional Chinese Medicine, Xiamen University Malaysia.

Cite this article: Jin KY, Kong YY, Aslam MS. A Narrative and Meta-Analytic Study of *in vivo* Efficiency of the Bioactive Compounds of Propolis in Tooth Decay. *Pharmacog Rev.* 2019;13(26):28-35.