

PHCOG REV. : Review Article

A Review: *Carissa Congesta*: Phytochemical Constituents, Traditional Use and Pharmacological properties

V. Devmurari*, Shivanand P, M.B. Goyani, S.Vaghani, N.P.Jivani.

Smt.R.B.Patel Mahila Pharmacy College, Kailash nagar, Bhavnagar Road, Atkot, Gujarat, India

*Author for corospondance:viraldev1985@gmail.com

Abstract

Various herbal drugs individually or in combination have been recommended for the treatment of different disease. The *C.congesta* commonly known as "karanda" has been recognized in different system of traditional medicine to cure various diseases. It contains several phytochemical constituents belonging to terpenoids category. The root is attributed with bitter, stomachic; anti-diarrhoeal, vermifuge and anti-anthelmintic properties. The unripe fruit is used medicinally as an astringent. The ripe fruit is taken as an antiscorbutic and remedy for biliousness. The leaf decoction is valued in cases of intermittent fever, diarrhoea, oral inflammation and earache. Additionally *C.congesta* has shown wide range of evidences for its cardiotoxic, hepatoprotective, free radical scavenging and xanthine oxidase inhibitory, histamine releasing, anti-rhumatic, antibacterial, antiviral and anticonvulsant activity. A higher gross heat value of this species indicates its higher potential to be used as good fuel source.

Key words: Karanda, Phytochemical constituents, Traditional uses, and Pharmacological properties.

Abbreviations:*C.carandas*=*Carissa congesta*, *B.subtillis*=*Bacillus subtillis*, *S.aureus*=*Staphylococcus aureus*, *S.fecalis*=*Streptococcus faecalis*, *E.coli*=*Escherichia coli*, *p.aeruginosa*=*Pseudomonas aeruginosa* and *S. typhimurium*=*Salmonella typhimurium*.

INTRODUCTION

Carissa congesta Wight (syn. *C. carandas* Auct., formerly widely shown as *C. carandas* L.) belong to *Apocynaceae*. It is called kerenda in Malaya, karaunda in India; Bengal currant or Christ's thorn in South India; namdaeng in Thailand; caramba, caranda, caraunda and perunkila in the Philippines (1). This species is a rank-growing, straggly, woody, climbing shrub, usually growing to 10 or 15 ft (3-5 m) high, sometimes ascending to the tops of tall trees; and rich in white, gummy latex. The branches, numerous and spreading, forming dense masses, are set with sharp thorns, simple or forked, up to 2 in (5 cm) long, in pairs in the axils of the leaves. The leaves are evergreen, opposite, oval or elliptic, 1 to 3 in (2.5-7.5 cm) long; dark-green, leathery, glossy on the upper surface, lighter green and dull on the underside. The fragrant flowers are tubular with 5 hairy lobes which are twisted to the left in the bud instead of to the right as in other species. They are white, often tinged with pink, and borne in terminal clusters of 2 to 12. The fruit, in clusters of 3 to 10, is oblong, broad-ovoid or round, 1/2 to 1 in (1.25-2.5 cm) long; has fairly thin but tough, purplish-red skin turning dark-purple or nearly black when ripe; smooth, glossy; enclosing very acid to fairly sweet, often bitter, juicy, red or pink, juicy pulp, exuding flecks of latex. There may be 2 to 8 small, flat, brown seeds. The karanda is native and common throughout much of India, Burma and Malacca and dry areas of Ceylon; is rather commonly cultivated in these areas as a hedge and for its fruit and the fruit is marketed in villages (2). The karanda was first fruited in the Philippines in 1915 and P.J. Wester described it in 1918 as "one of the best small fruits introduced into the Philippines within recent years." The present review will possibly act as bridge between traditional claim and modern therapy of *C.congesta*.



Figure 1 : Stem, fruit, flower, leaf and bark of *C.congesta*

TRADITIONAL USES

C.congesta is the best known as member of the genus as it has been used as a traditional medicinal plant over thousands of years in the ayurvedic system of medicine as it is practiced on the Indian sub continent. Thus, traditional uses of *C.congesta* are well established. The root is credited with bitter, stomachic, anti-diarrhoeal and anti-anthelmintic properties. The ripe fruits are utilized in curries, tarts, puddings and chutney. When only slightly under ripe, they are made into jelly. Green, sour fruits are made into pickles in India. With skin and seeds removed and seasoned with sugar and cloves, they have been popular as a substitute for apple in tarts. The unripe fruit is used medicinally as an astringent. The ripe fruit is taken as an antiscorbutic and remedy for biliousness. The fruits have been employed as agents in tanning and dyeing British residents in India undoubtedly favored the karanda as being reminiscent of gooseberries. Karanda leaves have furnished fodder for the tussar silkworm. The leaf decoction is valued in cases of intermittent fever, diarrhea, oral inflammation and earache. A paste of the pounded roots serves as a fly repellent. The root is employed as a bitter stomachic and vermifuge and it is an

ingredient in a remedy for itches. The roots contain salicylic acid and cardiac glycosides causing a slight decrease in blood pressure. The white or yellow wood is hard, smooth and useful for fashioning spoons, combs, household utensils and miscellaneous products of turnery. It is sometimes burned as fuel (3, 4).

PHYTOCHEMICAL CONSTITUENTS

The roots of *C. congesta* have yielded a number of volatile principles including 2-acetyl phenol (5, 6). Pal et al have reported a new lignan, carinol from root of *C. congesta* (7). Whereas studies carried out by Rastogi et al have led to isolation of a mixture of sesqui terpenes, namely carissone and carindone as a novel type of C₃₁ terpenoid (8-10). The leaves were reported to have triterpenoid constituents well as tannins, and a new isomer of urosolic acid namely carissic acid was also found (11, 12). Fruits of this plant were reported to contain a mixture of volatile constituents including 2-phenyl ethanol, linalool, β-caryophylline, isoamyl alcohol, benzyl acetate and a novel triterpenic alcohol, carissol. Enzymatic mild hydrolysis of polar glycoside from the plant yielded oderoside H, digitoxigenin and the sugars D-glucose and D-digitalose. Moreover, *C. congesta* contains crude protein 13%, polyphenols 7.8%, fixed oil 5.3% hydrocarbons 58% and free acid 31.4%. Higher gross heat values of this species indicate that it can be used as fuel source. Essential oil from *C. congesta* was found to contain coumarin (13).

It has been reported that fresh leaves of *C. congesta* contain four pentacyclic triterpenoids including one new constituent carissin and two hitherto unreported compounds. The structure of the new triterpenoid has been elucidated as 3-β-hydroxy-27-E-feruloyloxyurs-12-en-28-oic acid. Complete assignment of the protons of the compounds has also been made based on 2D NMR studies (14).




Figure 2 : Phytochemical constituents of *C. congesta*

PHARMACOLOGICAL ACTIVITIES

Anticonvulsant action

It has been reported that ethanolic extract of *C. congesta* has powerful anticonvulsant action on electrically and chemically induced seizures by unknown mode of action (15).

Cardiotonic activity

The alcoholic extract of roots of *C. congesta* exhibited cardiotonic activity and prolonged blood pressure lowering effect. An atorphous water-soluble polyglycoside possessing significant cardiac activity has been isolated. The cardiac activity of water-soluble fraction has been attributed to the presence of the glucosides of oderoside (16).

Free radicals scavenging and xanthine oxidase inhibitory activity

The free radical scavenging activity of ethanolic and aqueous

extracts of *C. congesta* been evaluated, in vitro, using 1,1-diphenyl-2-picrylhydrazyl (DPPH) radical scavenging method and compared with ascorbic acid, quercetin and apigenin. The plant has demonstrated good DPPH radical scavenging activity.

Inhibitory effects towards the in-vitro reaction of hypoxanthine and xanthine oxidase (XO) was also carried out in the presence of plant extract, aglycones quercetin, and apigenin along with allopurinol XO inhibitory activity of Plant extracts, was comparable with aglycones but less than allopurinol (17).

Potential Hepatoprotective and Antioxidant activity

It has been proved that *C. congesta* has hepatoprotective and antioxidant action by Chatterjee et al. They have used Wistar albino rats to estimate serum glutamate oxaloacetate transaminase (SGOT), serum glutamate pyruvate transaminase (SGPT), serum alkaline phosphatase (SALP), uric acid, total protein content and total bilirubin content. They have measured antioxidant action by estimating lipid peroxidation, reduced glutathione (GSH), super oxide dismutase (SOD) and catalase (CAT) activity in liver (18).

Histamine releasing activity

It has been proved that *C. congesta* have histamine releasing effect that was observed by estimating plasma histamine in cats, histamine content in rat hind-limb and histamine content of the lung tissue (19).

Antimicrobial action

It has been reported that ethanolic extract of *C. congesta* has potent antibacterial action against different test bacteria like *B. subtilis*, *S. aureus*, *S. faecalis*, *E. coli*, *P. aeruginosa* and *S. typhimurium*. Moreover ethanol extract has also showed considerable anticandidal action (20).

Antiviral action

It has been reported that ethanolic extract of *C. congesta* possess potent antiviral activity against Sindbis virus (SINV) at 3 µg/ml, polio virus (POLIO), at 6 µg/ml HIV-1, and herpes simplex virus (HSV) at 12 µg/ml (21).

It was known that Root of *C. congesta* had antirhumatic property (22). Isolated carissone derivatives exhibited an antizymotic, antibacterial and atropine like spasmolytic activity (23). The lipase activity of fruit has also been studied (24). An aqueous extract of the root demonstrated anthelmintic, spasmolytic, cardiotonic and hypertensive action in a preliminary pharmacological screening (25). It is reported that *C. congesta* has also good antipyretic activity (26).

CONCLUSION

Nowadays, herbal drugs are widely used as curative agent for different ailments. Concentrated extract of *C. congesta* can be found in various herbal preparations which are readily available in the market. *C. congesta* preparations have broad range of distribution in the market and employed by practitioner of natural health for the treatment of rheumatism. In the traditional system of medicine, *C. congesta* plant is used as cardiotonic, hepatoprotective, free radical scavenger and xanthine oxidase inhibiting agent, histamine releasing agent, antirhumatic, stomachic; antidiarrhoeal, vermifuge, antianthelmintic, astringent, antiscorbutic, antibacterial,

antiviral and anticonvulsant. Moreover, it has good potential to be used as potent energy source.

REFERENCES

1. S. K. Jain, *Dictionary of Indian Folk Medicines and Ethnobotany*, (Deep Publication, New Delhi, 1991).
2. K.R. Kirtikar, B.D. Basu. *Indian Medicinal Plants.*, Vol. II, Allahabad, 2003,1546- 1549.
3. G Addis, D. Abebe, K Urga . A survey of traditional medicinal plants in Shirka district, Arsi zone, Ethiopia. *Ethiopian Pharmaceutical Journal*. **19**: 30-47 (2001).
4. S.C. Pakrashi, S. Datta , PP. Ghosh-Dastidar. Indian medicinal plants- XVII.3 Phytochemical examination of *Carissa* SPP. *Phytochemistry* **7(3)**:495-6 (1968).
5. A.Zaki, S. El-Tohamy, S. El-Fattah. Study of Lipid content and volatile oil of the different organs of *Carissa carandas* Lin. and *Carissa grandiflora* Dc. growing in Egypt. *Egyptian Journal of Pharmaceutical Sciences* **22(1-4)**:127-41 (1983).
6. J. Pino, R. Marbot, C. Vazques . Volatile flavour constituents of Karnda (*Carissa carandas* L.) fruit. *Journal of Essential Oil Research*. **16(5)**: 432-4 (2004).
7. R. Pal, D.K. Kulshreshtha, R.P. Rastogi. A new lignan from *Carissa carandas*. *Phytochemistry*. **14(10)**: 2302-3 (1975).
8. B. Singh, R.P. Rastogi . The structure of carindone. *Phytochemistry*, **11(5)**:1797-801 (1972).
9. J. Reisch, R. Hussain, B. Krebs, M. Dartmann. The structure of carissone. *Monatshefte fuer Chemie* **121(11)**: 941-4 (1990).
10. D.V. Joshi, S. F. Boyce. Chemical Investigation of Root of *Carissa congesta*, Santapau. *Journal of Organic Chemistry*.**22**: 95-7 (1957).
11. S. Siddiqui, U. Ghani, S. Ali, S. Usmani, S. Begum. Triterpenoidal Constituents of the leaves of *Carissa carandas*. *Natural Products Research*. **17(3)**:153 (2003).
12. Z. Naim, M. Khan, S. Nizami. Isolation of a new isomer of ursolic acid from fruits and leaves of *Carissa carandas*. *Pakistan Journal of Scientific and Industrial Research*. **31(11)**: 753-5 (1988).
13. G. Chandra . Essential Oil of *Carissa carandas*. Examination of the benzene extract of the flowers and of the essential oil. *Soap, Perfumery & Cosmetics*. **45(6)**:551-6 (1972).
14. B.S. Siddiqui, U. Ghani U, S.T. Ali, S.B. Usmani, S. Begum. Triterpenoidal constituents of the leaves of *Carissa carandas*. *Nat Prod Res*. **17(3)**:153-8 (2003).
15. K. Hegde¹, S.P. Thakker², A.B. Joshi³, CS Shastri⁴, KS Chandrashekhar³, Anticonvulsant Activity of *Carissa carandas* Linn. Root Extract in Experimental Mice, *Tropical Journal of Pharmaceutical Research*. **8 (2)**: 117-125 (2009).
16. M.M Vohra, N. N. De. Comparative cardiotonic activity of *Carissa carandas* {L}. and *Carissa spinarum* {A}. *Indian journal of medical research*. **51(5)**: 937-940 (1963).
17. N.S. Ahmad^{1*}, M. Farman², M. H. Najmi¹, K.B. Mian¹and A. Hasan² Activity of polyphenolic plant extracts as scavengers of free radicals and inhibitors of xanthine oxidase. *Journal of Applied and Basic Science*. **2 (1)**: (2006).
18. M.L. Chatterjee, A.R. Roy. Pharmacological action of *Carissa carandas* root. *Bulletin of the Calcutta School of Tropical Medicine*, **13(1)**:14-6 (1965).
19. S.N. Joglekar and B.B. Gaitonde, Histamine releasing activity of *Carissa Carandas* roots (apocyanaceae), *Japanese. Journal of Pharmacy*. **20**: 367-372 (1970).
20. P. Jigna, N. Rathish, C. Sumitra. Preliminary screening of some folklore medicinal plants from western India for potential antimicrobial activity, *Indian J Pharmacol*, **37(6)**: 408-409 (2005).
21. S R.S.L. Tylor, Medicinal plants of Nepal: Ethnomedicine, Pharmacology and phytochemistry, Ph D Thesis, University of british Columbia, Sep 1996.
22. Parveen*, B. Upadhyay, S. Roy, A.Kumar¹, Traditional uses of medicinal plants among the rural communities of Churu district in the Thar Desert, India., *Journal of Ethnopharmacology* . **113**: 387–399 (2007).
23. A.P. Bhaduri, R.P. Rastogi, N.M. Khanna. Biologically active carissone derivatives. *Indian Journal of Chemistry*. **6**: 405 (1968)
24. V. Mala, Pahoja and M. Sethar, A review of enzymatic properties of lipase in plant, animals and microorganism. *Pakistan Journal of Applied Science*. **2(2)**: 474-484 (2002).
25. M.L. Chatterjee, A.R. Roy. Pharmacological action of *Carissa carandas* root. *Bulletin of the Calcutta School of Tropical Medicine*. **13(1)**:14-6 (1965).
26. PA Batugal, J Kanniah, S.Y. Lee and JT Oliver, *Medicinal Plants Research in Asia*, Volume I, (The frame work and project work plan, International Plant Genetic Resources Institute) 2004.